



A service of the National Library of Medicine
and the National Institutes of Health

My NCBI
[Sign In] [Register]
Entrez 2.0

[All Databases](#)
[PubMed](#)
[Nucleotide](#)
[Protein](#)
[Genome](#)
[Structure](#)
[OMIM](#)
[PMC](#)
[Journals](#)
[Books](#)

Search ☒ for

[Limits](#)
[Preview/Index](#)
[History](#)
[Clipboard](#)
[Details](#)
[About Entrez](#)
[Text Version](#)
[Entrez PubMed](#)
[Overview](#)
[Help | FAQ](#)
[Tutorials](#)
[New/Noteworthy](#)
[E-Utilities](#)
[PubMed Services](#)
[Journals Database](#)
[MeSH Database](#)
[Single Citation](#)
[Matcher](#)
[Batch Citation](#)
[Matcher](#)
[Clinical Queries](#)
[Special Queries](#)
[LinkOut](#)
[My NCBI](#)

- Search History will be lost after eight hours of inactivity.
- Search numbers may not be continuous; all searches are represented.
- To save search indefinitely, click query # and select Save in My NCBI.
- To combine searches use #search, e.g., #2 AND #3 or click query # for more options.

Search

Most Recent Queries

Time Result

[#5 Search Perez and CEA](#)

17:39:53 [19](#)
[#3 Search Freyre and CEA](#)

17:38:20 [2](#)
[#2 Search Tormo and CEA](#)

17:34:08 [1](#)
[#1 Search CB/ior-CEA.1](#)

17:33:28 [0](#)

Related Resources

[Order Documents](#)
[NLM Mobile](#)
[NLM Catalog](#)
[NLM Gateway](#)
[TOXNET](#)
[Consumer Health](#)
[Clinical Alerts](#)
[ClinicalTrials.gov](#)
[PubMed Central](#)
[Write to the Help Desk](#)
[NCBI | NLM | NIH](#)
[Department of Health & Human Services](#)
[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

WEST Search History

[Hide Items](#)[Restore](#)[Clear](#)[Cancel](#)

DATE: Thursday, May 31, 2007

Hide?	<u>Set Name</u>	<u>Query</u>	<u>Hit Count</u>
	<i>DB=PGPB,USPT,USOC,EPAB,JPAB,DWPI; PLUR=YES; OP=ADJ</i>		
<input type="checkbox"/>	L1	CB/ior-CEA.1	3

END OF SEARCH HISTORY

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSPTALAB1643

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS 1 Web Page for STN Seminar Schedule - N. America
NEWS 2 JAN 08 CHEMLIST enhanced with New Zealand Inventory of Chemicals
NEWS 3 JAN 16 CA/Caplus Company Name Thesaurus enhanced and reloaded
NEWS 4 JAN 16 IPC version 2007.01 thesaurus available on STN
NEWS 5 JAN 16 WPIDS/WPINDEX/WPIX enhanced with IPC 8 reclassification data
NEWS 6 JAN 22 CA/Caplus updated with revised CAS roles
NEWS 7 JAN 22 CA/Caplus enhanced with patent applications from India
NEWS 8 JAN 29 PHAR reloaded with new search and display fields
NEWS 9 JAN 29 CAS Registry Number crossover limit increased to 300,000 in
multiple databases
NEWS 10 FEB 15 PATDPASPC enhanced with Drug Approval numbers
NEWS 11 FEB 15 RUSSIAPAT enhanced with pre-1994 records
NEWS 12 FEB 23 KOREAPAT enhanced with IPC 8 features and functionality
NEWS 13 FEB 26 MEDLINE reloaded with enhancements
NEWS 14 FEB 26 EMBASE enhanced with Clinical Trial Number field
NEWS 15 FEB 26 TOXCENTER enhanced with reloaded MEDLINE
NEWS 16 FEB 26 IFICDB/IFIPAT/IFIUDB reloaded with enhancements
NEWS 17 FEB 26 CAS Registry Number crossover limit increased from 10,000
to 300,000 in multiple databases
NEWS 18 MAR 15 WPIDS/WPIX enhanced with new FRAGHITSTR display format
NEWS 19 MAR 16 CASREACT coverage extended
NEWS 20 MAR 20 MARPAT now updated daily
NEWS 21 MAR 22 LWPI reloaded
NEWS 22 MAR 30 RDISCLOSURE reloaded with enhancements
NEWS 23 APR 02 JICST-EPLUS removed from database clusters and STN
NEWS 24 APR 30 GENBANK reloaded and enhanced with Genome Project ID field
NEWS 25 APR 30 CHEMCATS enhanced with 1.2 million new records
NEWS 26 APR 30 CA/Caplus enhanced with 1870-1889 U.S. patent records
NEWS 27 APR 30 INPADOC replaced by INPADOCDB on STN
NEWS 28 MAY 01 New CAS web site launched
NEWS 29 MAY 08 CA/Caplus Indian patent publication number format defined
NEWS 30 MAY 14 RDISCLOSURE on STN Easy enhanced with new search and display
fields
NEWS 31 MAY 21 BIOSIS reloaded and enhanced with archival data
NEWS 32 MAY 21 TOXCENTER enhanced with BIOSIS reload
NEWS 33 MAY 21 CA/Caplus enhanced with additional kind codes for German
patents
NEWS 34 MAY 22 CA/Caplus enhanced with IPC reclassification in Japanese
patents

NEWS EXPRESS NOVEMBER 10 CURRENT WINDOWS VERSION IS V8.01c, CURRENT
MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP),
AND CURRENT DISCOVER FILE IS DATED 25 SEPTEMBER 2006.

NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS LOGIN Welcome Banner and News Items
NEWS IPC8 For general information regarding STN implementation of IPC 8

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 18:02:45 ON 31 MAY 2007

=> file caplus

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

0.21

0.21

FILE 'CAPLUS' ENTERED AT 18:02:59 ON 31 MAY 2007

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 31 May 2007 VOL 146 ISS 23

FILE LAST UPDATED: 30 May 2007 (20070530/ED)

Effective October 17, 2005, revised CAS Information Use Policies apply. They are available for your review at:

<http://www.cas.org/infopolicy.html>

=> s (CB-CEA-1)

13472 CB

4391 CBS

17515 CB

(CB OR CBS)

6551 CEA

196 CEAS

6725 CEA

(CEA OR CEAS)

9150913 1

L1 5 (CB-CEA-1)

(CB(W)CEA(W)1)

=> duplicate remove l1

PROCESSING COMPLETED FOR L1

L2 5 DUPLICATE REMOVE L1 (0 DUPLICATES REMOVED)

=> d l2 bib abs 1-5

L2 ANSWER 1 OF 5 CAPLUS COPYRIGHT 2007 ACS on STN

AN 1996:85276 CAPLUS

DN 124:172974

TI Bacterial expression and characterization of a modified scFv fragment from the anti-carcinoembryonic antigen monoclonal antibody CB-CEA.1
AU Vazquez, Javier E.; Perez, Lilia; Ayala, Martha; Canaan-Haden, Leonardo; De Lalla, C.; Sidoli, A.; Gavigondo, Jorge
CS Center Genetic Engineering and Biotechnology, Havana, 10600, Cuba
SO Biotecnologia Aplicada (1995), 12(2), 96-7
CODEN: BTAPEP; ISSN: 0864-4551
PB Sociedad Ibero-latinoamericana de Biotecnologia Aplicada a la Salud
DT Journal
LA English
AB The authors report here the cloning, bacterial expression (*Escherichia coli*), and mol. characterization of an anti-carcinoembryonic antigen (CEA) single-chain Fv antibody fragment with potential use in the localization of tumors in CEA-pos. carcinoma patients.

L2 ANSWER 2 OF 5 CAPLUS COPYRIGHT 2007 ACS on STN

AN 1993:78955 CAPLUS

DN 118:78955

TI Bacterial single-chain antibody fragments, specific for carcinoembryonic antigen

AU Ayala, M.; Duenas, M.; Santos, A.; Vazquez, J.; Menendez, A.; Silva, A.; Gavigondo, J. V.

CS Cent. Genet. Eng. Biotechnol., Cuba

SO BioTechniques (1992), 13(5), 790-2, 794-6, 798-9
CODEN: BTNQDO; ISSN: 0736-6205

DT Journal

LA English

AB Single-chain antibody (scFv) fragments were produced in bacteria specific for carcinoembryonic antigen (CEA). Polymerase chain reaction (PCR) was used for the cloning and modification of the heavy and light variable regions (VH and VL) of the mouse monoclonal antibody (MAb) CB-CEA.1. Culture supernatant, bacteria pellet and periplasm preps. were assayed in Western blot, and a protein of .apprx.27 kDa was identified with rabbit antibodies specific for the Fab of CB-CEA.1. Bacterial supernatant and periplasm preps. also inhibited the recognition of CEA by HRP-labeled CB-CEA.1 in ELISA. Periplasm preps. were purified by affinity chromatog. with specific anti-idiotypic monoclonal antibodies. The Western blot of the eluates identified a protein of .apprx.27 kDa that blocked the recognition of CEA by HRP-labeled CB-CEA.1 in ELISA.

L2 ANSWER 3 OF 5 CAPLUS COPYRIGHT 2007 ACS on STN

AN 1992:122152 CAPLUS

DN 116:122152

TI Primer design for the cloning of immunoglobulin heavy-chain leader-variable regions from mouse hybridoma cells using the PCR

AU Coloma, Maria J.; Larrick, James W.

CS Genelabs, Inc., Redwood City, CA, 94063, USA

SO BioTechniques (1991), 11(2), 152-4, 156

CODEN: BTNQDO; ISSN: 0736-6205

DT Journal

LA English

AB To facilitate the rapid cloning and sequencing of rearranged murine heavy-chain variable regions, a set of universal primers was designed using conserved sequences of leader (signal peptide), framework one and constant regions of the Ig heavy-chain genes. RNA was extracted from the mouse hybridoma cells secreting monoclonal antibodies: IOR-T3 (anti-CD3), C6 (anti-P1 of *N. meningitidis* B385), IOR-T1 (anti-CD6), CB-CEA.1 (anti-carcinoembryonic antigen), CB-Fib.1 (anti-human fibrin) and CB-Hep.2 (anti-hepatitis B surface antigen). First-strand cDNA was synthesized and amplified using PCR. The primers successfully amplified correct size fragments from cDNA prepared from all

hybridomas. These methods will facilitate the cloning and sequencing of mouse Ig variable regions.

L2 ANSWER 4 OF 5 CAPLUS COPYRIGHT 2007 ACS on STN
AN 1991:56891 CAPLUS
DN 114:56891
TI Specific amplification of rearranged immunoglobulin variable region genes from mouse hybridoma cells
AU Gavalondo-Cowley, Jorge V.; Coloma, Maria J.; Vazquez, Javier; Ayala, Marta; Macias, Amparo; Fry, Kirk E.; Larrick, James W.
CS Div. Hybridomas Anim. Models, Cent. Genet. Eng. Biotechnol., Havana, Cuba
SO Hybridoma (1990), 9(5), 407-17
CODEN: HYBRDY; ISSN: 0272-457X
DT Journal
LA English
AB This article describes how the polymerase chain reaction (PCR) and primers designed for conserved sequences of leader (L), framework one (FR1) and constant (CONST) regions of Ig light and heavy chain genes can be used for the cloning and sequencing of rearranged antibody variable regions from mouse hybridoma cells. RNA was extracted from the mouse hybridoma cells secreting MAbs: IOR-T3a (anti-CD3), C6 (anti-P1 of *Neisseria meningitidis* B385), IOR-T1 (anti-CD6), CB-CEA.1 (anti-carcinoembryonic antigen), and CB-Fib.1 (anti-human fibrin). First strand cDNA was synthesized and amplified using PCR. The newly designed primers are superior to others reported recently in the literature. Isolated PCR DNA fragments of C6 and IOR-T3a were sequenced after asym. amplification, or M13 cloning. The FR1/CONST primer combinations selectively amplified mouse light chains of groups kappa II, V, and VI, and heavy chains of groups IIa and IIc. The L/CONST primers for light chains amplified light chains from all 4 hybridomas. The methods greatly facilitate structural and functional studies of antibodies by reducing the efforts required to clone and sequence their variable regions.

L2 ANSWER 5 OF 5 CAPLUS COPYRIGHT 2007 ACS on STN
AN 1990:156288 CAPLUS
DN 112:156288
TI Relative positions of some epitopes on carcinoembryonic antigen
AU Larsson, Aake; Ghosh, Rahul; Hammarstroem, Sten
CS Dep. Immunol., Univ. Stockholm, Stockholm, S-10691, Swed.
SO Cancer Immunology Immunotherapy (1989), 30(2), 92-6
CODEN: CIIMDN; ISSN: 0340-7004
DT Journal
LA English
AB To investigate whether anti-(carcinoembryonic antigen) monoclonal antibodies (mAb) react with single or repeated epitopes, sandwich RIAs in homologous and heterologous combinations were performed. Four mAb (I-27, I-47, II-17, and to some degree II-16) gave homologous binding, while two mAb (I-38S1 and II-10) did not. All these mAb except II-10 react with repeated epitopes. The relative positions of the epitopes recognized by these mAb and of three addnl. mAb (II-6, II-7, and CB-CEA-1) were investigated using a plate antibody competition test with enzyme-labeled carcinoembryonic antigen (CEA). The mAb I-38S1, II-6, II-7, II-10, II-16, and CB-CEA-1 were mutually cross-reactive, and were classified as belonging to one epitope group. The mAb I-27 and I-47 fell outside this group and did not interfere with the binding of CEA conjugate to mAb II-17 either. They, therefore, represent a second epitope group. The mAb II-17 showed no interference with the binding of CEA to any of the other mAb and must therefore represent a third epitope group. The slopes of the plate antibody competition curves were used for calcn. of a correlation matrix, which in turn was used to depict the relative positions of the epitopes recognized by the mAb in the large group.

=> logoff

ALL L# QUERIES AND ANSWER SETS ARE DELETED AT LOGOFF

Welcome to DialogClassic Web(tm)

Dialog level 05.17.01D

Last logoff: 21may07 12:17:10

Logon file: 31may07 16:53:51

*** ANNOUNCEMENTS ***

NEW FILES RELEASED

***BIOSIS Previews Archive (File 552)

***BIOSIS Previews 1969-2007 (File 525)

***Engineering Index Backfile (File 988)

***Trademarkscan - South Korea (File 655)

RESUMED UPDATING

***File 141, Reader's Guide Abstracts

RELOADS COMPLETED

***Files 154 & 155, MEDLINE

***File 5, BIOSIS Previews - archival data added

***Files 340, 341 & 942, CLAIMS/U.S. Patents - 2006 reload now online

DATABASES REMOVED

Chemical Structure Searching now available in Prous Science Drug Data Report (F452), Prous Science Drugs of the Future (F453), IMS R&D Focus (F445/955), Pharmaprojects (F128/928), Beilstein Facts (F390), Derwent Chemistry Resource (F355) and Index Chemicus (File 302).

>>>For the latest news about Dialog products, services, content<<<

>>>and events, please visit What's New from Dialog at <<<

>>><http://www.dialog.com/whatsnew/>. You can find news about<<<

>>>a specific database by entering HELP NEWS <file number>.<<<

* * *

File 1:ERIC 1965-2007/Apr

(c) format only 2007 Dialog

Set Items Description

--- -----

Cost is in DialUnits

?

B 155, 159, 10, 203, 35, 5, 467, 73, 434, 34

31may07 16:54:34 User290558 Session D109.1

\$0.98 0.279 DialUnits File1

\$0.98 Estimated cost File1

\$0.18 INTERNET

\$1.16 Estimated cost this search

\$1.16 Estimated total session cost 0.279 DialUnits

SYSTEM:OS - DIALOG OneSearch

File 155:MEDLINE(R) 1950-2007/May 30

(c) format only 2007 Dialog

*File 155: Medline has been reloaded. Please see HELP NEWS 154 for information on 2007 changes.

File 159:Cancerlit 1975-2002/Oct

(c) format only 2002 Dialog

*File 159: Cancerlit is no longer updating. Please see HELP NEWS159.

File 10:AGRICOLA 70-2007/May

(c) format only 2007 Dialog

File 203:AGRIS 1974-2007/Mar
Dist by NAL, Intl Copr. All rights reserved
File 35:Dissertation Abs Online 1861-2007/May
(c) 2007 ProQuest Info&Learning
File 5:Biosis Previews(R) 1926-2007/May W4
(c) 2007 The Thomson Corporation
***File 5: BIOSIS has been enhanced with archival data. Please see
HELP NEWS 5 for information.**
File 467:ExtraMED(tm) 2000/Dec
(c) 2001 Informania Ltd.
File 73:EMBASE 1974-2007/May 24
(c) 2007 Elsevier B.V.
File 434:SciSearch(R) Cited Ref Sci 1974-1989/Dec
(c) 2006 The Thomson Corp
File 34:SciSearch(R) Cited Ref Sci 1990-2007/Jun W1
(c) 2007 The Thomson Corp

Set	Items	Description
-----	-------	-------------

---	-----	-----
-----	-------	-------

?

S	(CB (W) IOR (W) CEA (W) 1)
	442858 CB
	1497 IOR
	56221 CEA
	16945423 1
S1	0 (CB (W) IOR (W) CEA (W) 1)

?

S	(CEA AND MONOCLONAL)
	56221 CEA
	872021 MONOCLONAL
S2	8956 (CEA AND MONOCLONAL)

?

S	S2 AND CB
	8956 S2
	442858 CB
S3	115 S2 AND CB

?

S S3 AND (CEA (W) 1)
Processing
Processed 10 of 10 files ...
Completed processing all files
115 S3
56221 CEA
16945423 1
168 CEA(W)1
S4 17 S3 AND (CEA (W) 1)

?

RD S4
S5 5 RD S4 (unique items)

?

TYPE S5/FULL/1-5

5/9/1 (Item 1 from file: 155)
DIALOG(R) File 155:MEDLINE(R)

(c) format only 2007 Dialog. All rts. reserv.

09428616 PMID: 1418981

Bacterial single-chain antibody fragments, specific for carcinoembryonic antigen.

Ayala M; Duenas M; Santos A; Vazquez J; Menendez A; Silva A; Gavilondo J V

Division of Immunotechnology and Diagnostics, Center for Genetic Engineering and Biotechnology, La Habana, Cuba.

BioTechniques (UNITED STATES) Nov 1992, 13 (5) p790-9, ISSN 0736-6205--Print Journal Code: 8306785

Publishing Model Print

Document type: Journal Article

Languages: ENGLISH

Main Citation Owner: NLM

Record type: MEDLINE; Completed

Subfile: INDEX MEDICUS

We have produced single-chain antibody (scFv) fragments in bacteria specific for carcinoembryonic antigen (CEA). Polymerase chain reaction (PCR) was used for the cloning and modification of the heavy and light variable regions (VH and VL) of the mouse monoclonal antibody (MAb) CB-CEA.1. A 14-amino acid linker was used in the synthesis of the scFv gene. The VH and VL regions were amplified from cDNA by PCR using 5' end FR1 and 3' end constant region primers, and then sequenced. VH was then amplified by PCR using an exact 5' end FR1 primer, and a phosphorylated (PP) 3' end primer for J2 that also encoded the first 7 amino acids of the linker. VL was amplified with a PP 5' end primer for FR1, also encoding the remaining 7 amino acids of the linker, and a 3' end primer for J5, plus a stop codon and a BglII restriction site. The fragments were ligated and reamplified with the PP VH 5' and VL 3' end primers. The VH-linker-VL structure was blunt-cloned into expression vectors bearing the tryptophan promoter and pelB or ompA signal peptide sequences. Culture supernatant, bacteria pellet and periplasm preparations were assayed in Western blot and a protein of about 27 kDa was identified with rabbit antibodies specific for the Fab of CB-CEA.1. Bacterial supernatant and periplasm preparations also inhibited the recognition of CEA by HRP-labeled CB-CEA.1 in enzyme-linked immunosorbent assay (ELISA). Periplasm preparations were purified by affinity chromatography with specific anti-idiotypic MAbs. The Western blot of the eluates identified a protein of approximately 27 kDa that blocked the recognition of CEA by HRP-labeled CB-CEA.1 in ELISA. The VH-linker-VL structure was cloned into a vector bearing the lacZ promoter and the pelB signal peptide. The recombinant bacterial clones also expressed about 27 kDa scFv, specific for CEA.

Descriptors: *Antibodies, Neoplasm--immunology--IM; *Antibody Specificity --genetics--GE; *Carcinoembryonic Antigen--immunology--IM; *Immunoglobulin Fragments--immunology--IM; Amino Acid Sequence; Animals; Antibodies, Monoclonal--immunology--IM; Antibodies, Neoplasm--genetics--GE; Base Sequence; Chromatography, Affinity; Cloning, Molecular; DNA, Single-Stranded; Enzyme-Linked Immunosorbent Assay; Escherichia coli; Genetic Vectors; Hybridomas; Immunoglobulin Fragments--genetics--GE; Mice; Molecular Sequence Data; Plasmids; Polymerase Chain Reaction; Protein Binding; Recombinant Proteins--metabolism--ME

Molecular Sequence Databank No.: GENBANK/D10270; GENBANK/D10271; GENBANK/D10272; GENBANK/D10273; GENBANK/D10274; GENBANK/D10275; GENBANK/D10276; GENBANK/M63506; GENBANK/S49453; GENBANK/S49457

CAS Registry No.: 0 (Antibodies, Monoclonal); 0 (Antibodies, Neoplasm); 0 (Carcinoembryonic Antigen); 0 (DNA, Single-Stranded); 0 (Immunoglobulin Fragments); 0 (Recombinant Proteins)

Gene Symbol: scFv

Record Date Created: 19921210

Record Date Completed: 19921210

5/9/2 (Item 2 from file: 155)

DIALOG(R) File 155:MEDLINE(R)

(c) format only 2007 Dialog. All rts. reserv.

09010334 PMID: 1931008

Primer design for the cloning of immunoglobulin heavy-chain leader-variable regions from mouse hybridoma cells using the PCR.

Coloma M J; Larrick J W; Ayala M; Gavilondo-Cowley J V

Genelabs, Inc., Redwood City, CA 94063.

BioTechniques (UNITED STATES) Aug 1991, 11 (2) p152-4, 156, ISSN 0736-6205--Print Journal Code: 8306785

Publishing Model Print

Document type: Journal Article; Research Support, Non-U.S. Gov't

Languages: ENGLISH

Main Citation Owner: NLM

Record type: MEDLINE; Completed

Subfile: INDEX MEDICUS

To facilitate the rapid cloning and sequencing of rearranged murine heavy-chain variable regions, we have designed a set of universal primers using conserved sequences of leader (signal peptide), framework one and constant regions of the immunoglobulin heavy-chain genes. RNA was extracted from the mouse hybridoma cells secreting monoclonal antibodies: IOR-T3 (anti-CD3), C6 (anti-P1 of N. meningitidis B385), IOR-T1 (anti-CD6), CB-CEA.1 (anti-carcinoembryonic antigen), CB-Fib.1 (anti-human fibrin) and CB-Hep.2 (anti-hepatitis B surface antigen). First-strand cDNA was synthesized and amplified using PCR. The primers successfully amplified correct size fragments from cDNA prepared from all hybridomas. These methods will facilitate the cloning and sequencing of mouse immunoglobulin variable regions.

Descriptors: *Immunoglobulin Heavy Chains--genetics--GE; *Immunoglobulin Variable Region--genetics--GE; *Polymerase Chain Reaction; Animals; Base Sequence; Cell Line; Cloning, Molecular; DNA--chemical synthesis--CS; Hybridomas; Mice; Molecular Sequence Data; Polydeoxyribonucleotides--chemical synthesis--CS

CAS Registry No.: 0 (Immunoglobulin Heavy Chains); 0 (Immunoglobulin Variable Region); 0 (Polydeoxyribonucleotides); 9007-49-2 (DNA)

Record Date Created: 19911203

Record Date Completed: 19911203

5/9/3 (Item 3 from file: 155)

DIALOG(R) File 155:MEDLINE(R)

(c) format only 2007 Dialog. All rts. reserv.

08323758 PMID: 2611022

CEA in colonic adenocarcinomas and precancerous lesions. An immunohistochemical study with a novel monoclonal antibody.

Tormo B R; Gavilondo J V; Dominguez C; Freyre M; Rodriguez T; Biberfeld P

Instituto Nacional de Oncologia y Radiobiologia, Departamento de Biologia, La Habana, Cuba.

APMIS - acta pathologica, microbiologica, et immunologica Scandinavica (DENMARK) Dec 1989, 97 (12) p1073-80, ISSN 0903-4641--Print

Journal Code: 8803400

Publishing Model Print

Document type: Journal Article; Research Support, Non-U.S. Gov't

Languages: ENGLISH

Main Citation Owner: NLM

Record type: MEDLINE; Completed

Subfile: INDEX MEDICUS

Using a new anti-CEA (CB-CEA-1) murine monoclonal antibody, the expression of carcinoembryonic antigen (CEA) was studied in normal, premalignant and malignant human adult tissues with particular emphasis on colorectal mucosa. The CB-CEA-1 epitope was poorly expressed in normal adult tissues but was consistently found in colon cancers and adenomas in distinctive immunohistochemical patterns. Some apical staining was found with CB-CEA-1 in cells of normal colon mucosa whereas colon adenocarcinomas had a predominantly cytoplasmic staining pattern. Colonic adenomas presented a varied staining pattern. Some showed apical staining, others a CEA distribution pattern similar to that of adenocarcinomas, particularly those with a villous component. Our findings indicate a differential expression of CB-CEA-1 in adenoma cells in relation to their potential for malignant transformation. The possible usefulness of this Mab defined epitope for diagnostic and therapeutic purposes is indicated.

Descriptors: *Adenocarcinoma--metabolism--ME; *Carcinoembryonic Antigen--metabolism--ME; *Colonic Neoplasms--metabolism--ME; *Precancerous Conditions--metabolism--ME; Adenocarcinoma--pathology--PA; Antibodies, Monoclonal--diagnostic use--DU; Antibodies, Monoclonal--immunology--IM; Carcinoembryonic Antigen--immunology--IM; Colonic Neoplasms--diagnosis--DI; Colonic Neoplasms--pathology--PA; Humans; Immunohistochemistry; Precancerous Conditions--diagnosis--DI; Precancerous Conditions--pathology--PA

CAS Registry No.: 0 (Antibodies, Monoclonal); 0 (Carcinoembryonic Antigen)

Record Date Created: 19900302

Record Date Completed: 19900302

5/9/4 (Item 4 from file: 155)

DIALOG(R)File 155:MEDLINE(R)

(c) format only 2007 Dialog. All rts. reserv.

08300833 PMID: 2480845

Relative positions of some epitopes on carcinoembryonic antigen.

Larsson A; Ghosh R; Hammarstrom S

Department of Immunology, University of Stockholm, Sweden.

Cancer immunology, immunotherapy - CII (GERMANY, WEST) 1989, 30 (2)
p92-6, ISSN 0340-7004--Print Journal Code: 8605732

Publishing Model Print

Document type: Journal Article; Research Support, Non-U.S. Gov't

Languages: ENGLISH

Main Citation Owner: NLM

Record type: MEDLINE; Completed

Subfile: INDEX MEDICUS

To investigate whether anti-(carcinoembryonic antigen) monoclonal antibodies (mAb) react with single or repeated epitopes, sandwich radioimmunoassays in homologous and heterologous combinations were performed. Four mAb (I-27, I-47, II-17 and to some degree II-16) gave homologous binding while two mAb (I-38S1 and II-10) did not. Taken together with previous immunoprecipitation studies we conclude that all these mAb except II-10 react with repeated epitopes. The relative positions of the epitopes recognized by these mAb and of three additional mAb (II-6, II-7 and CB-CEA-1) were investigated using a plate antibody competition test with enzyme-labelled carcinoembryonic antigen (CEA). mAb I-38S1, II-6, II-7, II-10, II-16 and CB-CEA-1 were mutually cross-reactive, and were classified as belonging to one epitope group. mAb I-27 and I-47 fell outside this group and did not interfere with the binding of CEA conjugate to mAb II-17 either. They therefore represent a second epitope group. mAb

II-17 showed no interference with the binding of CEA to any of the other mAb and must therefore represent a third epitope group. The slopes of the plate antibody competition curves were used for calculation of a correlation matrix, which in turn was used to depict the relative positions of the epitopes recognized by the mAb in the large group.

Descriptors: *Carcinoembryonic Antigen--immunology--IM; *Epitopes --analysis--AN; Antibodies, Monoclonal--diagnostic use--DU; Humans

CAS Registry No.: 0 (Antibodies, Monoclonal); 0 (Carcinoembryonic Antigen); 0 (Epitopes)

Record Date Created: 19900129

Record Date Completed: 19900129

5/9/5 (Item 1 from file: 5)

DIALOG(R)File 5:Biosis Previews(R)

(c) 2007 The Thomson Corporation. All rts. reserv.

13167253 BIOSIS NO.: 199698635086

Bacterial expression and characterization of a modified scFv fragment from the anti-carcinoembryonic antigen monoclonal antibody CB.CEA.1

AUTHOR: Vazquez Javier E (Reprint); Perez Lilia (Reprint); Ayala Martha (Reprint); Canaan-Haden Leonardo (Reprint); De Lalla C; Sidoli A; Gavilondo Jorge (Reprint)

AUTHOR ADDRESS: Cent. Genet. Eng. Biotechnol., P.O. Box 6162, Havana 10600, Cuba**Cuba

JOURNAL: Biotecnologia Aplicada 12 (2): p96-97 1995 1995

DOCUMENT TYPE: Article

RECORD TYPE: Citation

LANGUAGE: English

DESCRIPTORS:

MAJOR CONCEPTS: Biochemistry and Molecular Biophysics; Infection;

Metabolism; Oncology--Human Medicine, Medical Sciences; Physiology

BIOSYSTEMATIC NAMES: Hominidae--Primates, Mammalia, Vertebrata, Chordata, Animalia

ORGANISMS: human (Hominidae)

COMMON TAXONOMIC TERMS: Animals; Chordates; Humans; Mammals; Primates; Vertebrates

MISCELLANEOUS TERMS: ANTIBODY RECOGNITION PATTERN; CARCINOEMBRYONIC

ANTIGEN; MOLECULAR CHARACTERIZATION; NOTE; TUMOR LOCALIZATION

CONCEPT CODES:

10064 Biochemistry studies - Proteins, peptides and amino acids

10068 Biochemistry studies - Carbohydrates

10506 Biophysics - Molecular properties and macromolecules

13004 Metabolism - Carbohydrates

13012 Metabolism - Proteins, peptides and amino acids

24006 Neoplasms - Biochemistry

31000 Physiology and biochemistry of bacteria

36002 Medical and clinical microbiology - Bacteriology

BIOSYSTEMATIC CODES:

86215 Hominidae

?

S (CB (W) CEA (W) 1)

442858 CB

56221 CEA

16945423 1

S6 20 (CB (W) CEA (W) 1)

?

S S6 AND (THERAP? OR TREAT?)

Processing

Processed 10 of 10 files ...

Completed processing all files

20 S6
9076329 THERAP?
10091335 TREAT?
S7 4 S6 AND (THERAP? OR TREAT?)

?

RD S7

S8 1 RD S7 (unique items)

?

TYPE S8/FULL/1

8/9/1 (Item 1 from file: 155)

DIALOG(R) File 155:MEDLINE(R)

(c) format only 2007 Dialog. All rts. reserv.

08323758 PMID: 2611022

CEA in colonic adenocarcinomas and precancerous lesions. An immunohistochemical study with a novel monoclonal antibody.

Tormo B R; Gaviñondo J V; Dominguez C; Freyre M; Rodriguez T; Biberfeld P
Instituto Nacional de Oncología y Radiobiología, Departamento de Biología,
La Habana, Cuba.

APMIS - acta pathologica, microbiologica, et immunologica Scandinavica (DENMARK) Dec 1989, 97 (12) p1073-80, ISSN 0903-4641--Print

Journal Code: 8803400

Publishing Model Print

Document type: Journal Article; Research Support, Non-U.S. Gov't

Languages: ENGLISH

Main Citation Owner: NLM

Record type: MEDLINE; Completed

Subfile: INDEX MEDICUS

Using a new anti-CEA (CB-CEA-1) murine monoclonal antibody, the expression of carcinoembryonic antigen (CEA) was studied in normal, premalignant and malignant human adult tissues with particular emphasis on colorectal mucosa. The CB-CEA-1 epitope was poorly expressed in normal adult tissues but was consistently found in colon cancers and adenomas in distinctive immunohistochemical patterns. Some apical staining was found with CB-CEA-1 in cells of normal colon mucosa whereas colon adenocarcinomas had a predominantly cytoplasmic staining pattern. Colonic adenomas presented a varied staining pattern. Some showed apical staining, others a CEA distribution pattern similar to that of adenocarcinomas, particularly those with a villous component. Our findings indicate a differential expression of CB-CEA-1 in adenoma cells in relation to their potential for malignant transformation. The possible usefulness of this Mab defined epitope for diagnostic and therapeutic purposes is indicated.

Descriptors: *Adenocarcinoma--metabolism--ME; *Carcinoembryonic Antigen--metabolism--ME; *Colonic Neoplasms--metabolism--ME; *Precancerous Conditions--metabolism--ME; Adenocarcinoma--pathology--PA; Antibodies, Monoclonal--diagnostic use--DU; Antibodies, Monoclonal--immunology--IM; Carcinoembryonic Antigen--immunology--IM; Colonic Neoplasms--diagnosis--DI; Colonic Neoplasms--pathology--PA; Humans; Immunohistochemistry; Precancerous Conditions--diagnosis--DI; Precancerous Conditions--pathology--PA

CAS Registry No.: 0 (Antibodies, Monoclonal); 0 (Carcinoembryonic Antigen)

Record Date Created: 19900302

Record Date Completed: 19900302

?

Set	Items	Description
S1	0	(CB (W) IOR (W) CEA (W) 1)
S2	8956	(CEA AND MONOCLONAL)
S3	115	S2 AND CB
S4	17	S3 AND (CEA (W) 1)
S5	5	RD S4 (unique items)
S6	20	(CB (W) CEA (W) 1)
S7	4	S6 AND (THERAP? OR TREAT?)
S8	1	RD S7 (unique items)

?